arXiv:0912.5501v2 [math.NT] 5 Jan 2010

THE NUMBER OF POINTS ON AN ELLIPTIC CURVE WITH SQUARE x-COORDINATES

YU TSUMURA

Abstract. Let K be a finite field. We know that half of the elements of K^* are square. So it is natural to ask how many of them appear as x-coordinates of points on an elliptic curve over K. We consider a specific class of elliptic curves over finite fields and show that half of the x-coordinates on an elliptic curve are square.

1. Introduction.

Let K be a finite field of characteristic different from 2. Let $a, b \in K$ satisfy $b \neq 0$ and $r = a^2 - 4b \neq 0$. We consider a class of elliptic curves over K given by $E : y^2 = x^3 + ax^2 + bx$.

We know that half of the elements of K^* are square. So the natural question is how many points on an elliptic curve have square x-coordinates. Since half of the elements of K^* are square, we expect that half of the points on $E(K)$ have square x-coordinate.

We show that this is the case when the elliptic curve is $E : y^2 = x^3 + ax^2 + bx$.

2. Result

The main result is the following.

Theorem 2.1. *Let $E(K)$ be an elliptic curve defined above over a finite field K. Let $S = \{P \in E(K) | P = (x, y)$ with x being square or $P = \infty\}$, where ∞ is the identity of $E(K)$. Then we have*

$$\#S = \begin{cases} \#E(K)/2 & \text{if } b \text{ is a square in } K \\ \#E(K)/2 + 1 & \text{if } b \text{ is not a square in } K. \end{cases}$$

Proof. Consider $E' : Y^2 = X^3 - 2aX^2 + rX$. Then there is an isogeny $\phi : E' \longrightarrow E$ of degree 2 defined by

$$\phi : (X, Y) \longmapsto \left(\frac{Y^2}{4X^2}, \frac{Y(r - X^2)}{8X^2} \right). \tag{2.1}$$

(See page 70, Example 4.5 in [1].)

2010 *Mathematics Subject Classification.* Primary 14H52; Secondary 11G20.

Then by the isomorphism theorem we have, $E'(K)/\mathrm{Ker}(\phi) \cong \mathrm{Im}(\phi)$. Since E and E' are isogenous, it is easy to see that $\#E(K) = \#E'(K)$. (See page 153, Exercise 5.4 (a) in [1].) Also we have $\mathrm{Ker}(\phi) = \{\infty, (0,0)\}$. Hence we have

$$\#\mathrm{Im}(\phi) = \#E(K)/2. \tag{2.2}$$

Now we show that

$$S = \begin{cases} \mathrm{Im}(\phi) & \text{if } b \text{ is a square in } K \\ \mathrm{Im}(\phi) \cup \{(0,0)\} & \text{if } b \text{ is not a square in } K. \end{cases}$$

Let R denote the right hand side. It is easy to see that the preimage of $(0,0)$ by ϕ over the algebraic closure $\bar{K}$ is $\{(a \pm 2\sqrt{b}, 0)\}$. Hence the preimage is in $E(K)$ if and only if b is a square in K. So $(0,0)$ is in both S and R.

Suppose $P \in S$. If P is ∞, then clearly $P \in \mathrm{Im}(\phi)$. If $P = (x,y) \neq (0,0)$ with x square, then we need to solve

$$\frac{Y^2}{4X^2} = x, \frac{Y(r - X^2)}{8X^2} = y$$

for $(X,Y) \in E'(K)$. By the first equation, using the relation $Y^2 = X^3 - 2aX^2 + rX$, we have $X^2 - 2(a+2x)X + r = 0$. Hence we get $X = a + 2x \pm 2\sqrt{x^2 + ax + b}$. Now since $y^2 = x(x^2 + ax + b)$ and x is a nonzero square, we see that $x^2 + ax + b$ is also a square. Hence $X \in K$. By the second equation for Y, we solve $Y = 8yX^2(r - X^2)^{-1} \in K$. Here $r - X^2 \neq 0$ since $y \neq 0$. Hence we get a point $(X,Y) \in E'(K)$ such that $\phi(X,Y) = (x,y)$. Therefore, we have $S \subset R$.

The converse follows by (2.1) and we have $R = S$.

Now we have

$$\#S = \begin{cases} \#\mathrm{Im}(\phi) & \text{if } b \text{ is a square in } K \\ \#\mathrm{Im}(\phi) + 1 & \text{if } b \text{ is not a square in } K. \end{cases}$$

By the equation 2.2,

$$\#S = \begin{cases} \#E(K)/2 & \text{if } b \text{ is a square in } K \\ \#E(K)/2 + 1 & \text{if } b \text{ is not a square in } K. \end{cases}$$

□

References

1. Joseph H. Silverman, *The arithmetic of elliptic curves*, second ed., Graduate Texts in Mathematics, vol. 106, Springer, Dordrecht, 2009. MR MR2514094

Department of Mathematics, Purdue University 150 North University Street, West Lafayette, Indiana 47907-2067

E-mail address: `ytsumura@math.purdue.edu`